CARDIOLOGY

1. **In complex therapy of *cardiovascular pathology*,** use **the standard course of immunomodulation, hepatoprotection, reparation and adaptation with ERBISOL® medication**

A course of therapy for 20 days 30 ampoules of ERBISOL® (e)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
6^{00} - 9^{00}				e	e	e	e	e	e	e	e	e	e							
21^{00} - 24^{00}	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	↕3 days↕			10 days										↕	7 days					↕

2. **In complex therapy of *ischemic heart disease with chronic heart failure of blood circulation of functional class I-II according NYHA classification*,** use **the intensive course of immunomodulation, reparation hepato-, cardiovascular protection with ERBISOL® Extra medication**

A course of therapy for 21 days 60 ampoules of ERBISOL® Extra (2+)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
6^{00} – 9^{00}		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21^{00} - 24^{00}	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	↕1↕							19 days													↕1↕

3. **In complex therapy of *ischemic heart disease with chronic heart failure of blood circulation of functional class III according NYHA* classification*,** use **the standard prolonged course of ERBISOL® Extra medication**

A course of therapy for 20-30 days 40-60 ampoules of ERBISOL® Extra (+)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6^{00} – 9^{00}	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
21^{00} – 24^{00}	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	↕	20-30 days																												↕

* **In patients with more severe chronic heart failure of blood circulation, this standard prolonged course is appropriate only after its stabilization at the level of NYHA functional class III.**

4. **In complex therapy of *combined pathology of the cardiovascular system, on the background of which type 2 diabetes mellitus developed,*** use **the systemic therapy of intensive combined course of immunomodulation, reparation, hepato-, cardiovascular protection and adaptation with ERBISOL® and ERBISOL®Extra medications, and intensive prolonged course of immunomodulation, reparation, and hepato-, cardiovascular protection with ERBISOL®Extra medication.**

The 1st intensive combined course of therapy for 21 days
20 ampoules of ERBISOL® (e) + 40 ampoules of ERBISOL® Extra (2+)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
6^{00} - 9^{00}		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21^{00} - 24^{00}	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	

↕1↕ 19 days ↕1↕

2 months after the end of the 1st course, begin the 2nd course of therapy.

The 2nd intensive prolonged course of therapy for 21-31 days
60-90 ampoules of ERBISOL® Extra (+)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
6^{00} - 9^{00}		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21^{00} - 24^{00}	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	

↕1↕ 19-**29** days ↕1↕

5. In complex therapy of patients with *chronic cardiovascular pathology (in postinfarction or postinsult periods),* use the standard prolonged course of ERBISOL® Extra medication

A course of therapy for 20-30 days 40-60 ampoules of ERBISOL® Extra (+)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6^{00} - 9^{00}	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
21^{00} - 24^{00}	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

↕ 20-**30** days ↕

Medications should preferably be injected on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:

e - 1 (one) ampoule per 2 ml of ERBISOL® intramuscularly;

+ - 1 (one) ampoule per 2 ml of ERBISOL® Extra intramuscularly.