

ENDOCRINOLOGY

1. In complex therapy *of type 1 and type 2 diabetes* it is recommended to use **the standard course** of immunomodulation, hepatoprotection, repairation and adaptation with ERBISOL[®] medication

A course of therapy for 20 days 30 ampoules of ERBISOL[®] (e)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Time																				
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e	e	e	e	e	e							
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	3 days↓			10 days							↓	7 days							↓	

This course is recommended to be held regularly 3-4 times a year.

2. In complex therapy **of type 1 and type 2 diabetes** it is recommended to use **the standard semi-prolonged course** of immunomodulation, repairation, and hepato-, cardiovascular protection with ERBISOL[®] Extra medication

A course of therapy for 30 days 50 ampoules of ERBISOL[®] Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30		
Time																																
9 ⁰⁰ - 11 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+		
21 ⁰⁰ - 24 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
	↓	20 days																			↓	10 days										↓

This course is recommended to be held regularly 2-3 times a year.

3. In complex therapy *of type 1 and type 2 diabetes with cardiovascular complications* it is recommended to use **the standard prolonged course** with ERBISOL[®] Extra medication

A course of therapy for 20-30 days 40-60 ampoules of ERBISOL[®] Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Time																														
6 ⁰⁰ - 9 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
21 ⁰⁰ - 24 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	↓	20-30 days																												↓

This course is recommended to be held regularly 1-2 times a year.

4. In complex therapy *of type 2 diabetes against the background of the combined pathology of the cardiovascular system*, it is recommended to use **the intensive prolonged course of ERBISOL[®] Extra**

A course of therapy for 21-31 days 60-90 ampoules of ERBISOL[®] Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	
Time																																
6 ⁰⁰ - 9 ⁰⁰		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	
21 ⁰⁰ - 24 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
		19-29 days																														
	↓1↓																												↓1↓			

Medications should preferably be injected on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:

e - 1 (one) ampoule per 2 ml of ERBISOL[®] intramuscularly;

+ - 1 (one) ampoule per 2 ml of ERBISOL[®] Extra intramuscularly;