GYNECOLOGY

In complex treatment of infertility caused by persistent infection, ovarian cysts and leiomyomas, use systemic therapy of pulsation and standard courses of immunomodulation, hepatoprotection, reparation and adaptation with ERBISOL® medication.

The 1st pulsation course of therapy for 30 days
50 ampoules of ERBISOL® (e)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
$6^{\underline{00}} - 9^{\underline{00}}$				e	e	e	e	e			e	e	e	e	e			e	e	e	e	e			e	e	e	e	e	
$21^{\underline{00}} - 24^{\underline{00}}$	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e

3 days ↕ 5 days ↕2days↕ 5 days ↕2days↕ 5 days ↕2days↕ 5 days ↕1↕

2 months after the end of the 1st course, begin the 2nd course of therapy.

The 2nd standard course of therapy for 20 days
30 ampoules of ERBISOL® (e)

Days / Time	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$6^{\underline{00}} - 9^{\underline{00}}$				e	e	e	e	e	e	e	e	e	e							
$21^{\underline{00}} - 24^{\underline{00}}$	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e

↕3 days↕ 10 days ↕ 7 days ↕

Injections of medications should preferably be administered on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:
e - 1 (one) per 2 ml of ERBISOL® intramuscularly.