

Infectious diseases

1. In complex therapy of *infectious diseases of bacterial etiology*, use **the standard course** of immunomodulation, hepatoprotection, reparation and adaptation with ERBISOL® medication.

**A course of therapy for 20 days
30 ampoules of ERBISOL® (e)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
Time																					
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e	e	e	e	e	e								
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	↓3 days↑			10 days							↓7 days↑										

2. In complex therapy of *diseases caused by intracellular pathogens*, use **the standard course** of immunotherapy, and hepatoprotection with ERBISOL® ULTRapharm medication.

**A course of therapy for 22 days
40 ampoules of ERBISOL® ULTRapharm (u)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
Time																						
6 ⁰⁰ - 9 ⁰⁰				u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u
21 ⁰⁰ - 24 ⁰⁰	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u
	↓3 days↑			18 days																		↓1↑

If necessary, the course of treatment can be continued up to 33 days

3. In complex therapy of *persistent infections*, use **the systemic therapy of pulsation and standard courses** of ERBISOL® medication.

**The 1st pulsation course of therapy for 30 days
50 ampoules of 2 ml ERBISOL® (e)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Time																															
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e			e	e	e	e	e			e	e	e	e	e			e	e	e	e	e		
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	
	3 days↓			5 days			↓2days↑		5 days					↓2days↑		5 days			↓2days↑		5 days			↓1↑							

2 months after the end of the 1st course, begin the 2nd course of therapy.

**The 2nd standard course of therapy for 20 days
30 ampoules of ERBISOL® (e)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
Time																					
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e	e	e	e	e	e								
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	↓3 days↑			10 days							↓7 days↑										

4. In complex therapy of *severe forms of persistent infections*, use **the systemic therapy with standard prolonged course** of ERBISOL® ULTRApHarm medication and **standard course** of ERBISOL® medication.

The 1st standard prolonged course of therapy for 30 days
50 ampoules of ERBISOL® ULTRApHarm (u)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Time																															
6 ⁰⁰ - 9 ⁰⁰				u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u								
21 ⁰⁰ - 24 ⁰⁰	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	
	↓3days↑			20 days												↓			7 days			↓									

After the end of the 1st course, begin the 2nd course without interruption.

The 2nd standard course of therapy for 20 days
30 ampoules of ERBISOL® (e)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
Time																					
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e	e	e	e	e									
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	↓3 days↑			10 days								↓			7 days			↓			

5. In complex therapy of *recurrent and mixed forms of persistent infections*, use **systemic therapy with intensive course** of ERBISOL® ULTRApHarm medication and with **standard course** of ERBISOL® medication.

The 1st intensive course of therapy for 21 day
50 ampoules of ERBISOL® ULTRApHarm (u)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Time																					
6 ⁰⁰ - 9 ⁰⁰				u	u	2u	2u	2u	2u	2u	2u	2u	2u	2u	2u	2u	2u	u	u	u	
21 ⁰⁰ - 24 ⁰⁰	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u
	↓ 3+2 days ↓					12 days										↓3+1days↓					

Immediately after the end of the 1st course, the 2nd standard course of ERBISOL® medication should be began.

The 2nd standard course of therapy for 20 days
30 ampoules of ERBISOL® (e)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
Time																					
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e	e	e	e	e									
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	↓3 days↑			10 days								↓			7 days			↓			

Medications should preferably be injected on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:

- e - 1 (one) ampoule per 2 ml of ERBISOL® intramuscularly;
- + - 1 (one) ampoule per 2 ml of ERBISOL® Extra intramuscularly;
- u - 1 (one) ampoule per 2 ml of ERBISOL® ULTRApHarm intramuscularly.