

NEUROLOGY

- In complex therapy *for rehabilitation of postinsult states*, use **the systemic prolonged course** of immunomodulation, reparation, and hepato-, cardiovascular protection with ERBISOL[®] Extra medication.

A course of therapy for 20-30 days
40-60 ampoules of ERBISOL[®] Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Time																															
6 ⁰⁰ - 9 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
21 ⁰⁰ - 24 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	↓																													↓	
		20-30 days																													

- In complex therapy *for rehabilitation of patients with demyelinating polyneuropathies, Parkinson's disease*, use **semi-intensive of pulsation course** with ERBISOL[®] Extra medication.

A course of therapy with 2-3 five-days for 25-30 days
with 2 five-days for 25 days 60 ampoules of ERBISOL[®] Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25			
Time																												
6 ⁰⁰ - 9 ⁰⁰	+	+	+	+	+	+	+	2+	2+	2+	2+	2+	+	+	2+	2+	2+	2+	2+	+	+	+	+	+	+			
21 ⁰⁰ - 24 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+			
	↓	7-9 days					↓	5 days					↓	2days		↓	5 days					↓	6 days					↓

with 3 five-days for 30 days 75 ampoules of ERBISOL[®] Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30				
Time																																		
6 ⁰⁰ - 9 ⁰⁰	+	+	+	+	+	+	+	2+	2+	2+	2+	2+	+	+	2+	2+	2+	2+	2+	+	+	2+	2+	2+	2+	2+	+	+	+	+				
21 ⁰⁰ - 24 ⁰⁰	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				
	↓	7-9 days					↓	5 days					↓	2days		↓	5 days					↓	5 days					↓	4 days					↓

- In complex therapy *of multiple sclerosis* use **intensive combined pulsation course** of immunotherapy, reparation and hepato-, cardiovascular protection with ERBISOL[®] ULTRApHarm and ERBISOL[®] Extra.

A course of therapy for 24 days

20 ampoules of ERBISOL[®] ULTRApHarm (u) + 35-40 ampoules of ERBISOL[®] Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24								
Time																																
6 ⁰⁰ - 9 ⁰⁰		+	+	2+	2+	2+	2+	2+	+	+	2+	2+	2+	2+	2+	+	+	2+	2+	2+	2+	2+										
21 ⁰⁰ - 24 ⁰⁰	u	u	u	u	u	u	u	u	+	+	u	u	u	u	u	+	+	u	u	u	u	u	u	u								
		3 days			↓	5 days					↓	2days		↓	5 days					↓	2days		↓	5 days					↓	2days		↓

Muscle tension resulting in the stiffness of the patient's movement can be reduced by the dose of ERBISOL[®] Extra in the 2nd and 3rd five-day periods and by use of the medication alternately in the morning - one day 2 ampoules **2+** (4.0 ml), another 1 ampoule **2+** (2.0 ml).

Note:

- + - 1 (one) ampoule per 2 ml of ERBISOL[®] Extra intramuscularly;
- u - 1 (one) ampoule per 2 ml of ERBISOL[®] ULTRApHarm intramuscularly.