

# RHEUMATOLOGY

1. In complex therapy *of seasonal exacerbations of osteoarthritis*, use **the intensive course** of immunomodulation, reparation and hepato-, cardiovascular protection with ERBISOL<sup>®</sup> Extra medication

### A course of therapy for 21 days

60 ampoules of ERBISOL<sup>®</sup> Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
Time																						
6 <sup>00</sup> - 9 <sup>00</sup>		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21 <sup>00</sup> - 24 <sup>00</sup>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

↑1↓
19 days
↑1↓

2. In complex therapy *of mix arthropathias*, use **the semi-intensive course** of ERBISOL<sup>®</sup> Extra medication.

### A course of therapy for 23 days

60 ampoules of ERBISOL<sup>®</sup> Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
Time																							
6 <sup>00</sup> - 9 <sup>00</sup>		+	+	+	+	+	+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21 <sup>00</sup> - 24 <sup>00</sup>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

↑1↓
6 days
↓
15 days
↑1↓

3. In complex therapy *of exacerbation of rheumatoid arthritis* against the background of the basic "gold standard" therapy with methotrexate, use **the intensive combined course** of immunotherapy, reparation and hepato-, cardiovascular protection with ERBISOL<sup>®</sup> ULTRApHarm and ERBISOL<sup>®</sup> Extra medications

### A course of therapy for 21 day

20 ampoules of ERBISOL<sup>®</sup> ULTRApHarm (u) + 40 ampoules of ERBISOL<sup>®</sup> Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
Time																						
6 <sup>00</sup> - 9 <sup>00</sup>		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21 <sup>00</sup> - 24 <sup>00</sup>	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u	u

↑1↓
19 days
↑1↓

4. In complex therapy *of rheumatoid arthritis with side effects* from the effect of basic "gold standard" therapy with methotrexate, use **the semi-intensive course** of immunomodulation, reparation and hepato-, cardiovascular protection with ERBISOL<sup>®</sup> Extra medication.

### A course of therapy for 23 days

60 ampoules of ERBISOL<sup>®</sup> Extra (+)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
Time																							
6 <sup>00</sup> - 9 <sup>00</sup>		+	+	+	+	+	+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21 <sup>00</sup> - 24 <sup>00</sup>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

↑1↓
6 days
↓
15 days
↑1↓

Medications should preferably be injected on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:

+ - 1 (one) ampoule per 2 ml of ERBISOL<sup>®</sup> Extra intramuscularly.

u - 1 (one) ampoule per 2 ml of ERBISOL<sup>®</sup> ULTRApHarm intramuscularly.