

# SURGERY

1. In the *post-operation period*, use **the intensive combined prolonged course** of immunomodulation, repairation, hepato-, cardiovascular protection and adaptation with ERBISOL<sup>®</sup> and ERBISOL<sup>®</sup> Extra medications.

### A course of therapy for 27 days

**30 ampoules of ERBISOL<sup>®</sup> (e) + 40 ampoules of ERBISOL<sup>®</sup> Extra (2+)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	
Time																												
6 <sup>00</sup> - 9 <sup>00</sup>					e	e	e	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21 <sup>00</sup> - 24 <sup>00</sup>	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	↓	7 days						↓	20 days																			↓

2. In complex therapy of *furunculosis and trophic ulcers*, use **systemic therapy of pulsation and intensive combined course** of immunomodulation, repairation, hepato-, cardiovascular protection and adaptation with ERBISOL<sup>®</sup> and ERBISOL<sup>®</sup> Extra medications.

### The 1<sup>st</sup> pulsation course of therapy for 30 days

**50 ampoules of ERBISOL<sup>®</sup> (e)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Time																															
6 <sup>00</sup> - 9 <sup>00</sup>				e	e	e	e	e			e	e	e	e	e			e	e	e	e	e			e	e	e	e	e		
21 <sup>00</sup> - 24 <sup>00</sup>	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	
	↓	3 days		↓	5 days			↓	2 days		↓	5 days			↓	2 days		↓	5 days			↓	2 days		↓	5 days			↓	1	↓

2 months after the end of the 1<sup>st</sup> course or from the moment of recurrence, begin the 2<sup>nd</sup> course of therapy

### The 2<sup>nd</sup> intensive combined course of therapy for 21 day

**20 ampoules of ERBISOL<sup>®</sup> (e) + 40 ampoules of ERBISOL<sup>®</sup> Extra (2+)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Time																					
6 <sup>00</sup> - 9 <sup>00</sup>		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+
21 <sup>00</sup> - 24 <sup>00</sup>	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	↓	19 days																			↓

Medications should preferably be injected on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:

e - 1 (one) ampoule per 2 ml of ERBISOL<sup>®</sup> intramuscularly;

+ - 1 (one) ampoule per 2 ml of ERBISOL<sup>®</sup> Extra intramuscularly.