

TOXICOLOGY

1. In complex therapy *of mild forms of poisoning*, use **the standard course** of immunomodulation, hepatoprotection, reparation and adaptation with ERBISOL[®] medication

A course of therapy for 20 days 30 ampoules of ERBISOL[®] (e)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
Time																					
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e	e	e	e	e	e								
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
	3 days ↓			10 days							↓	7 days			↓						

2. In complex therapy **of severe forms of poisoning**, use **pulsation course** of ERBISOL[®] medication

A course of therapy for 30 days 50 ampoules of ERBISOL[®] (e)

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Time																															
6 ⁰⁰ - 9 ⁰⁰				e	e	e	e	e			e	e	e	e	e			e	e	e	e	e			e	e	e	e	e		
21 ⁰⁰ - 24 ⁰⁰	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	
	3 days ↓			5 days			↓	2days ↓		5 days			↓	2days ↓		5 days			↓	2days ↓		5 days			↓	1 ↓					

Medications should preferably be injected on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:

e - 1 (one) ampoule per 2 ml of ERBISOL[®] intramuscularly.