

# TRAUMATOLOGY AND ORTHOPAEDY

In complex therapy *of severe injuries and fractures*, use **the systemic therapy of pulsation and intensive combined courses** of immunomodulation, reparation, hepato-, cardiovascular protection and adaptation with ERBISOL<sup>®</sup> Extra and ERBISOL<sup>®</sup> medications.

**The 1<sup>st</sup> pulsation course of therapy for 30 days**  
**50 ampoules of ERBISOL<sup>®</sup> (e)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Time																														
6 <sup>00</sup> - 9 <sup>00</sup>				e	e	e	e	e			e	e	e	e	e			e	e	e	e	e			e	e	e	e	e	
21 <sup>00</sup> - 24 <sup>00</sup>	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e
3 days ↓			5 days ↓			↑2days↓		5 days ↓			↑2days↓		5 days ↓			↑2days↓		5 days ↓			↑2days↓		5 days ↓			↑1↓				

2 months after the end of the 1<sup>st</sup> course, begin the 2<sup>nd</sup> course of therapy

**The 2<sup>nd</sup> intensive combined course of therapy for 21 day**  
**20 ampoules of ERBISOL<sup>®</sup> (e) + 40 ampoules of ERBISOL<sup>®</sup> Extra (2+)**

Days	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
Time																						
6 <sup>00</sup> - 9 <sup>00</sup>		2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	2+	
21 <sup>00</sup> - 24 <sup>00</sup>	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	e	
↑1↓		19 days																			↑1↓	

Medications should preferably be injected on an empty stomach 1-2 hrs before or 2-3 hrs after meals. Tea, juices, water, etc. can be consumed at any time.

Note:

e - 1 (one) ampoule per 2 ml of ERBISOL<sup>®</sup> intramuscularly;

+ - 1 (one) ampoule per 2 ml of ERBISOL<sup>®</sup> Extra intramuscularly.